

Entanglement and Quantum Phase Transition in Low Dimensional Spin Systems

Yan Chen¹, Paolo Zanardi², Z. D. Wang^{1,3}, and F. C. Zhang¹

¹*Department of Physics and Center of Theoretical and Computational Physics,
University of Hong Kong, Pokfulam Road, Hong Kong, China*

²*Institute for Scientific Interchange Foundation, Viale Settimio Severo 65, I-10133 Torino, Italy*

³*National Laboratory of Solid State Microstructures, Nanjing University, Nanjing, China*

(Dated: February 1, 2008)

Entanglement of the ground states in XXZ and dimerized Heisenberg spin chains as well as in a two-leg spin ladder is analyzed by using the spin-spin concurrence and the entanglement entropy between a selected sublattice of spins and the rest of the system. In particular, we reveal that quantum phase transition points/boundaries may be identified based on the analysis on the local extreme of this entanglement entropy, which is illustrated to be superior over the concurrence scenario and may enable us to explore quantum phase transitions in many other systems including higher dimensional ones.

PACS numbers: 03.67.Mn, 05.70.Jk, 75.10.Jm

Introduction. Quantum entanglement has attracted a lot of attention for its potential applications in quantum information processing [1, 2]. More recently, entanglement has also been recognized to play an important role in the study of quantum many particle physics, and experimental measurements have demonstrated that it can affect the macroscopic properties of solids [3]. There have been a number of theoretical studies of the entanglement and quantum phase transitions [4] in one dimensional spin systems and in interacting fermion and boson systems [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]. These studies showed that the entanglement can exhibit a non-trivial behavior in condensed matter systems, such as the coincidence of the singularity in the entanglement and quantum phase transition point in certain systems [17]. Most of the previous works were mainly focused on spin chains and the spin-spin concurrence [1] was used to describe (two-particle) the entanglement. The scaling behavior of entanglement between a block of contiguous spins and the rest of the system in a spin chain has been studied both near and at the quantum critical point [18], and a connection with quantum phase transitions was elaborated.

In this paper, we adopt a preferable and distinct way to partition the system to study the ground state of three low dimensional spin systems: the XXZ spin chain, dimerized Heisenberg chain, and two-leg spin ladder. We find that the entanglement entropy is superior over concurrence in revealing quantum transition points. In the dimerized Heisenberg system, this entanglement entropy has maxima and minima which are in one-to-one correspondence with the transition points, while the concurrence fails to locate them. Even though the spin ladder system exhibits a complex phase structure, our entanglement entropy scenario is likely to enable us to identify the phase boundaries. It is expected that our approach is more efficient and powerful in exploring quantum phase transitions in various systems.

The models. We here study the entanglement in three types of spin-1/2 systems: XXZ chain, dimerized Heisen-

berg chain, and two-leg XXZ ladder, with the Hamiltonian given by

$$H_{XXZ} = J \sum_{i=1}^N (\mathbf{S}_i^x \mathbf{S}_{i+1}^x + \mathbf{S}_i^y \mathbf{S}_{i+1}^y + \Delta \mathbf{S}_i^z \mathbf{S}_{i+1}^z), \quad (1)$$

$$H_D = \sum_{i=1}^{N/2} (J_1 \mathbf{S}_{2i-1} \cdot \mathbf{S}_{2i} + J_2 \mathbf{S}_{2i} \cdot \mathbf{S}_{2i+1}), \quad (2)$$

$$H_L = J \sum_{i=1}^2 \sum_{j=1}^N (\mathbf{S}_{i,j}^x \mathbf{S}_{i,j+1}^x + \mathbf{S}_{i,j}^y \mathbf{S}_{i,j+1}^y + \Delta \mathbf{S}_{i,j}^z \mathbf{S}_{i,j+1}^z) + J' \sum_{j=1}^N (\mathbf{S}_{1,j}^x \mathbf{S}_{2,j}^x + \mathbf{S}_{1,j}^y \mathbf{S}_{2,j}^y + \Delta \mathbf{S}_{1,j}^z \mathbf{S}_{2,j}^z), \quad (3)$$

where $\mathbf{S}_i$ is the spin operator, N is the linear dimension of the lattice (we adopt periodic boundary conditions). The anisotropy parameter is denoted by Δ in the XXZ model. J_1 and J_2 are two nearest-neighbor exchange couplings in the dimerized Heisenberg Hamiltonian. In the ladder system, the Hamiltonian contains intersite interactions along the chains (J) and the rungs (J').

To quantify the entanglement in these spin systems, we first adopt concurrence to describe two-spin entanglement [1]. This quantity can be calculated from correlation functions $G^{\alpha\alpha} := \langle \sigma^\alpha \otimes \sigma^\alpha \rangle$, ($\alpha = x, y, z$) [6]:

$$C = \frac{1}{2} \max \left[0, 2|G^{xx} + G^{yy}| - G^{zz} - 1 \right], \quad (4)$$

where the two site indexes are omitted. Since the correlation function decays rapidly, the concurrence is nonzero only between two closer sites. Therefore, the concurrence seems to provide limited information on the quantum phase transition. In a different approach, Vidal *et al.* [18] studied the entanglement between a block of L contiguous spins B_L and the rest of the system R_L . The entropy of entanglement for the ground state Ψ_g is given by,

$$S_L := -\text{tr}(\rho_L \log_2 \rho_L), \quad (5)$$




FIG. 1: The ground state concurrence (a) and the entanglement entropy per site S_L/L (b) in terms of Δ in XXZ spin chain for various lattice sizes ($N = 2L$). The inset of (b) displays the sublattice bipartition.

where $\rho_L \equiv \text{tr}_{B_L} |\Psi_g\rangle\langle\Psi_g|$ is the reduced density matrix for B_L , a block of L spins. The scaling of the entanglement in the block size L may allow one to establish a connection between S_L and the quantum phase transition: $S_L \sim \ln L$ at the critical point, and $S_L \sim \text{const}$ away from the critical point; while this idea is applicable to few one dimensional models. Here we adopt a distinct way that bi-partitions the system into two sublattices B_L and R_L , with the largest number of connecting bonds between them [19, 20]. The motivation of such partition is to best reveal the correlation between the two sublattices. We will see that in most cases the considered entanglement entropy between the two subsystems scales linearly with the size of the *boundary* between them [notice that in the case [18], the boundary is zero-dimensional i.e., one single spin][21].

All the above three Hamiltonians have a rotational symmetry around the z -axes. The calculations presented below are carried out in an invariant subspace with $S_z = 0$. Lanczos algorithms are employed to calculate the ground state $|\Psi_g\rangle$, from which we construct the density matrix for the whole system. We then obtain the reduced density matrix ρ_L by tracing out R_L , and compute its Von Neumann entropy in Eq.(5).

XXZ-chain. Let us first consider the one-dimensional XXZ model. As a simple toy model, a great deal of work has been devoted to analyze its entanglement and quantum phase transition [22]. It is well known that $\Delta = 1$ is the antiferromagnetic isotropic point, $\Delta = -1$ is the

ferromagnetic isotropic point, and $\Delta = 0$ is the pure XY point. This allows us to describe the various domains as a function of Δ . The system is in an Ising ferromagnetic phase at $\Delta < -1$, Ising antiferromagnetic phase at $\Delta > 1$, and XY phase at $-1 < \Delta < 1$. In Fig. 1(a), we show the pairwise concurrence as a function of anisotropy Δ for different number of sites L . Because the concurrence here is expressed in terms of the two-site correlation function of the nearest-neighbor sites, its value converges quickly as L increases. As we can see, the concurrence at one phase transition point $\Delta = 1$ reaches the maximum while the concurrence emerges at another transition point $\Delta = -1$ as L approaches infinity. These two quantum phase transition points can be identified from the analysis of concurrence in this case. We now address the entanglement entropy of the sublattice $B_L = \{\text{odd} - \text{sites}\}$ with $L = N/2$ [23]. Fig. 1(b) displays the numerical results of the entanglement entropy S_L/L as a function of Δ . We obtain the reduced subsystem of crosses by tracing out the spin degree of freedom on circle points. As the system is in the vicinity of the quantum phase transition point, we may expect S_L/L to reach its extreme value. We find that the transition points $\Delta = 1$ and $\Delta = -1$ correspond to the maximum and minimum of the sublattice entanglement entropy. As for the latter case, a simple analysis of the scaling behavior around the minimum point shows that the location of the transition point approaches to $\Delta = -1$ and $S_L/L = 0$ as the size of the subsystem increases. Thus at $L \rightarrow \infty$, $S_L/L \rightarrow 0$ for $\Delta < -1$. Therefore, the two transition points can be clearly specified and a distinct connection between quantum phase transition and the entanglement entropy has been established.

Dimerized Heisenberg chain. We now consider the dimerized Heisenberg chain [24]; this model is characterized by an alternation of strong and weak bonds between two nearest neighboring spins. In the case of $0 < J_2/J_1 \ll 1$, the ground state is just an ensemble of $N/2$ uncoupled dimers around the strong bonds. Consequently, there is an energy gap of order of J_1 to separate the singlet ground state ($S_z = 0$) from the first excited state with $S_z = 1$. All the spins are locked into singlet states. At $J_2/J_1 = 1$, the system is reduced to the isotropic antiferromagnetic Heisenberg chain that is quasi-long-range ordered, and belongs to a different universality class from the dimerized system. In the case of $J_2/J_1 < 0$, the ground state is a product of spin singlet pairs coupled by ferromagnetic bonds. We can define two concurrences C_1 and C_2 , which correspond to two nearest-neighboring spins coupled by bonds J_1 and J_2 , respectively. The pairwise concurrence as a function of J_2/J_1 for different lattice size L is shown in Fig. 2(a). It can be easily seen that the value of concurrence is independent of the lattice size. In the limit of strong dimerization, $J_2/J_1 = 0$ (or equivalently $J_2/J_1 \rightarrow \infty$), the concurrence C_1 (C_2) reaches a maximum value. However, the concurrence analysis does not enable us to identify the transition point $J_2/J_1 = 1$ unambiguously. Below,




FIG. 2: For various lattice sizes N in a dimerized Heisenberg chain, (a) the ground state concurrences C_1 (solid line) and C_2 (dotted line) versus Δ , and (b) S_L/L as a function of the sublattice size L . The inset in (b) shows the divergence of the $d^2(S_L/L)/d(J_2/J_1)^2$ at the block entanglement minimum point as the system size increases.

we examine the entanglement entropy of the chosen odd-site sublattice (or the even-site sublattice). In this case, the two transition points are found to correspond to a local maximum and a local minimum in the entanglement entropy [see Fig. 2(b)]. Here it is interesting to notice that the second derivative of S_L/L with respect to J_2/J_1 seems to exhibit singularity at the minimum point $J_2/J_1 = 1$ (see Fig. 2(b) inset), which corresponds to the only gapless point. Due to the limitation of our computational resources, a detailed scaling analysis is not available. In addition, it is seen that S_L/L is weakly size dependent and converges quickly to a fixed value at the transition point with increasing lattice size.

Spin ladders. Spin ladders have been attracting great interest in the field of low dimensional spin systems in recent several years [25]. The physics of Heisenberg ladders can be understood in some special limits. For large and negative J' , the 2-leg ladder is equivalent to an $S = 1$ spin chain, and the system is gapful [26]. For large and positive J' , the two spins along the rung are locked into a singlet state. The ground state is a collection of spin singlets on each rung and is gapful. The phase diagram of two coupled XXZ chains was previously determined by utilizing Hamiltonian mappings and Abelian bosonization [27]. Previous studies presumably indicated five distinct phases in this complicated phase diagram as a function of Δ and J'/J . We first examine the dependence on the coupling constant of two nearest-neighbor spin concurrence: this analysis failed to describe the rich




FIG. 3: S_L/L ($L = 8$) as a function of the J'/J and Δ in a coupled 2-leg XXZ spin ladder system. The results for $L = 6$ are essentially same.

structure of the phase diagram. Below we report our study on the entanglement entropy of the chosen subsystem and its relevance to quantum phase transition point. The spatial profiles and contour plots of S_L/L as a function of J'/J and Δ are displayed in Fig. 3. From our analysis from above-mentioned two models, we know that both the ridges and valleys may correspond to possible phase boundaries. For example, one can distinguish the boundary $J'/J = 0$ (ridge or valley), $\Delta = 1$ for $J'/J < 0$ (ridge), $\Delta = -1$ for $J' < 0$ (ridge). Derived from the numerical results, Fig. 4 shows the schematic phase boundaries of a coupled 2-leg XXZ spin ladder system (the dashed line denotes the ridge of its derivative). Most phase boundaries are in agreement with those in Ref. 26. Our results suggest the existence of a new quantum phase in the region $\Delta < -1.5$ and $J'/J > 0.5$, which has not been reported in previous studies [27, 28, 29].

Finally, it is important to emphasize that, in comparison with the previous investigations [18], we do not have to deal with a large system i.e., a large L , to capture its scaling behavior. This indicates that the finite size effects are not important in our approach; numerical results for a quite small system may disclose relevant information about the quantum phase transition points. In this sense, the present approach is broadly applicable to many other systems including spin and electron systems in higher dimension [19], not limited merely to the spin-chains.

Summary. We have investigated the ground-state entanglement in three one-dimensional spin systems by using both two-spin concurrence and entanglement entropy of the preferably chosen sublattice. Concurrence may provide some partial insights about certain quantum phase transition points. However, our entanglement entropy scenario allows us to establish a distinct connection between its local maxima/minima and transition




FIG. 4: Schematic phase boundaries of a coupled 2-leg XXZ spin ladder system.

points, which is promising for shedding a new light on the understanding of quantum phase transition and quantum entanglement.

We are grateful to Y. Q. Li, T. K. Lee, F. Mila and X. Q. Wang for useful discussions. This work was supported by the RGC grants of Hong Kong, the HKU URC Seed Funding, and the NSFC under Grant No. 10429401.

-
- [1] W.K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
 - [2] M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
 - [3] S. Ghosh, *et al.*, Nature **425**, 48 (2003).
 - [4] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, England, 2000).
 - [5] K.M. O'Connor and W.K. Wootters, Phys. Rev. A **63**, 052302 (2001).
 - [6] P. Zanardi, Phys. Rev. A **65**, 042101(2002).
 - [7] X. Wang, and P. Zanardi, Phys. Lett. A **301**, 1(2002).
 - [8] L.F. Santos, Phys. Rev. A **67**, 062306 (2003).
 - [9] S.J. Gu, H.Q. Lin, and Y.Q. Li, Phys. Rev. A **68**, 042330 (2003).
 - [10] F. Verstraete, M.A. Martin-Delgado, and J.I. Cirac, Phys. Rev. Lett. **92**, 087201 (2004).
 - [11] J.K. Pachos and M.B. Plenio, Phys. Rev. Lett. **93**, 056402 (2004).
 - [12] J.J. Garcia-Ripoll, M.A. Martin-Delgado, and J.I. Cirac, quant-ph/0404566.
 - [13] T.C. Wei, *et al.*, quant-ph/0405162.
 - [14] M.F. Yang, Phys. Rev. A **71**, 030302 (2005).
 - [15] S.J. Gu, S.S. Deng, Y.Q. Li, and H.Q. Lin, Phys. Rev. Lett. **93**, 086402 (2004).
 - [16] P. Giorda, and P. Zanardi, quant-ph/0311058.
 - [17] A. Osterloh, *et al.*, Nature **416**, 608 (2002).
 - [18] G. Vidal, J.I. Latorre, E. Rico, and A. Kitaev, Phys. Rev. Lett, **90**, 227902 (2003).
 - [19] Y. Chen, Z.D. Wang, F.C. Zhang (unpublished).
 - [20] This entanglement entropy satisfies the subadditivity property, see for example, A. Wehrl, Rev. Mod. Phys. **50**, 221 (1978).
 - [21] M.B. Plenio, *et al.*, quant-ph/0405142; A. Hamma, R. Ionicioiu, and P. Zanardi, quant-ph/0406202.
 - [22] T. Giamarchi, *Quantum Physics in One Dimension* (Oxford Science Publications, Oxford, 2004)
 - [23] We have examined the entanglement between the subsystem of L contiguous spins and its environment, and found that such a partition does not provide clear connection with the quantum critical point.
 - [24] M. Hase, I. Terasaki, and K. Uchinokura, Phys. Rev. Lett. **70**, 3651 (1993).
 - [25] E. Dagotto and T.M. Rice, Science, **271**, 618 (1996).
 - [26] F.D.M. Haldane, Phys. Rev. Lett. **60**, 635 (1988).
 - [27] S.P. Strong and A.J. Millis, Phys. Rev. Lett. **69**, 2419 (1992).
 - [28] E. Orignac and T. Giamarchi, Phys. Rev. B, **57**, 5812 (1998).
 - [29] It is quite clear each chain is in Ising ferromagnetic phase at $\Delta \ll -1$. For large and positive J' , the two spins along the rung are locked into singlet state. So this new phase may have two ferromagnetic chains with spin singlet pair along the rung direction.